


Fysioterapi til behandling af hofteartrose

FYSIOTERAPEUT, PH.D. LONE RAMER MIKKELSEN

1. Flachs E, Eriksen L, Koch M. Sygdomsbyrden i Danmark – sygdomme. Sundhedsstyrelsen. 2015.
 2. Cross M, Smith E, Hoy D, Nolte S, Ackerman I, Fransen M, et al. The global burden of hip and knee osteoarthritis: estimates from the global burden of disease 2010 study. *Ann Rheum Dis* 2014 Jul;73(7):1323-1330.
 3. Guillemin F, Rat AC, Mazieres B, Pouchot J, Fautrel B, Euller- Ziegler L, et al. Prevalence of symptomatic hip and knee osteoarthritis: a two-phase population-based survey. *Osteoarthritis Cartilage* 2011 Nov;19(11):1314-1322.
 4. Bartels EM, Juhl CB, Christensen R, Hagen KB, Danneskiold-Samsoe B, Dagfinrud H, et al. Aquatic exercise for the treatment of knee and hip osteoarthritis. *Cochrane Database Syst Rev* 2016 Mar 23;3:CD005523.
 5. Nelson AE, Allen KD, Golightly YM, Goode AP, Jordan JM. A systematic review of recommendations and guidelines for the management of osteoarthritis: The chronic osteoarthritis management initiative of the U.S. bone and joint initiative. *Semin Arthritis Rheum* 2014 Jun;43(6):701-712.
 6. Brand CA, Ackerman IN, Bohensky MA, Bennell KL. Chronic disease management: a review of current performance across quality of care domains and opportunities for improving osteoarthritis care. *Rheum Dis Clin North Am* 2013 Feb;39(1):123-143.
 7. Judd DL, Thomas AC, Dayton MR, Stevens-Lapsley JE. Strength and functional deficits in individuals with hip osteoarthritis compared to healthy, older adults. *Disabil Rehabil* 2014;36(4):307-312.
 8. Fautrel B, Hilliquin P, Rozenberg S, Allaert FA, Coste P, Leclerc A, et al. Impact of osteoarthritis: results of a nationwide survey of 10,000 patients consulting for OA. *Joint Bone Spine* 2005 May;72(3):235-240.
 9. Fransen M, McConnell S, Hernandez-Molina G, Reichenbach S. Exercise for osteoarthritis of the hip. *Cochrane Database Syst Rev* 2014 Apr 22;(4):CD007912. doi(4):CD007912.
 10. Thomee R. A comprehensive treatment approach for patellofemoral pain syndrome in young women. *Phys Ther* 1997 Dec;77(12):1690-1703.
 11. Sundhedsstyrelsen. National klinisk retningslinje for hofteartrose. Ikke-kirurgisk behandling og genoptræning efter total hoftealloplastik 2016.
 12. Hermann A, Holsgaard-Larsen A, Zerahn B, Mejdahl S, Overgaard S. Preoperative progressive explosive-type resistance training is feasible and effective in patients with hip osteoarthritis scheduled for total hip arthroplasty - a randomized controlled trial. *Osteoarthritis Cartilage* 2016 Jan;24(1):91-98.
 13. Bieler T, Siersma V, Magnusson SP, Kjaer M, Christensen HE, Beyer N. In hip osteoarthritis, Nordic Walking is superior to strength training and home-based exercise for improving function. *Scand J Med Sci Sports* 2016 Apr 30.
 14. Foley A, Halbert J, Hewitt T, Crotty M. Does hydrotherapy improve strength and physical function in patients with osteoarthritis - a randomised controlled trial comparing a gym based and a hydrotherapy based strengthening programme. *Ann Rheum Dis* 2003 Dec;62(12):1162-1167.
 15. Teirlinck CH, Luijsterburg PA, Dekker J, Bohnen AM, Verhaar JA, Koopmanschap MA, et al. Effectiveness of exercise therapy added to general practitioner care in patients with hip osteoarthritis: a pragmatic randomized controlled trial. *Osteoarthritis Cartilage* 2016 Jan;24(1):82-90.
-



16. French HP, Cusack T, Brennan A, Caffrey A, Conroy R, Cuddy V, et al. Exercise and manual physiotherapy arthritis research trial (EMPART) for osteoarthritis of the hip: a multicenter randomized controlled trial. *Arch Phys Med Rehabil* 2013 Feb;94(2):302-314.
 17. Villadsen A, Overgaard S, Holsgaard-Larsen A, Christensen R, Roos EM. Immediate efficacy of neuromuscular exercise in patients with severe osteoarthritis of the hip or knee: a secondary analysis from a randomized controlled trial. *J Rheumatol* 2014 Jul;41(7):1385-1394.
 18. Oosting E, Jans MP, Dronkers JJ, Naber RH, Dronkers-Landman CM, Appelman-de Vries SM, et al. Preoperative home-based physical therapy versus usual care to improve functional health of frail older adults scheduled for elective total hip arthroplasty: a pilot randomized controlled trial. *Arch Phys Med Rehabil* 2012 Apr;93(4):610-616.
 19. Skou ST. Træning som smertestillende medicin ved artrose er effektiv og uden alvorlige bivirkninger *Ugeskr Læger* 2016 Jun 13;178(24):V68364.
 20. Skou ST, Derosche CA, Andersen MM, Rathleff MS, Simonsen O. Nonoperative treatment improves pain irrespective of radiographic severity. A cohort study of 1,414 patients with knee osteoarthritis. *Acta Orthop* 2015;86(5):599-604.
 21. Juhl C, Christensen R, Roos EM, Zhang W, Lund H. Impact of exercise type and dose on pain and disability in knee osteoarthritis: a systematic review and meta-regression analysis of randomized controlled trials. *Arthritis Rheumatol* 2014 Mar;66(3):622-636.
 22. Abbott JH, Robertson MC, Chapple C, Pinto D, Wright AA, Leon de la Barra S, et al. Manual therapy, exercise therapy, or both, in addition to usual care, for osteoarthritis of the hip or knee: a randomized controlled trial. 1: clinical effectiveness. *Osteoarthritis Cartilage* 2013 Apr;21(4):525-534.
 23. Poulsen E, Hartvigsen J, Christensen HW, Roos EM, Vach W, Overgaard S. Patient education with or without manual therapy compared to a control group in patients with osteoarthritis of the hip. A proof-of-principle three-arm parallel group randomized clinical trial. *Osteoarthritis Cartilage* 2013 Oct;21(10):1494-1503.
 24. Skou ST, Roos EM. GLA:D® Annual Report 2015. 2016.
-