


Fysioterapi til mennesker med demens

Fuld referenceliste

- (1) Nationalt Videnscenter for Demens.
- (2) Sundheds- og Ældreministeriet (2016): Statusrapport på demensområdet i Danmark, http://www.sum.dk/~media/Filer%20-%20Publikationer_i_pdf/2016/Statusrapport-demens-2016/Statusrapport-paa-demensomraadet-i-dk.ashx.
- (3) Groot C, Hooghiemstra AM, Raijmakers PG, Berckel BN, Scheltens P, Scherder EJ, et al. The Effect of Physical Activity on Cognitive Function in Patients with Dementia: A Meta-Analysis of Randomized Control Trials. *Ageing research reviews*. 2015.
- (4) Bruce-Keller AJ, Brouillette RM, Tudor-Locke C, Foil HC, Cahia WP, Nye DM, et al. Relationship between cognitive domains, physical performance, and gait in elderly and demented subjects. *J Alzheimers Dis*. 2012;30:899-908.
- (5) Eggermont LH, Gavett BE, Volkens KM, Blankevoort CG, Scherder EJ, Jefferson AL, et al. Lower-extremity function in cognitively healthy aging, mild cognitive impairment, and Alzheimer's disease. *Archives of physical medicine and rehabilitation*. 2010;91:584-8.
- (6) Suttanon P, Hill KD, Said CM, Logiudice D, Lautenschlager NT, Dodd KJ. Balance and mobility dysfunction and falls risk in older people with mild to moderate Alzheimer disease. *American journal of physical medicine & rehabilitation / Association of Academic Physiatrists*. 2012;91:12-23.
- (7) Ijmker T, Lamoth CJ. Gait and cognition: the relationship between gait stability and variability with executive function in persons with and without dementia. *Gait & posture*. 2012;35:126-30.
- (8) Scherder E, Eggermont L, Swaab D, van Heuvelen M, Kamsma Y, de Greef M, et al. Gait in ageing and associated dementias; its relationship with cognition. *Neuroscience and biobehavioral reviews*. 2007;31:485-97.
- (9) Vidoni ED, Billinger SA, Lee C, Hamilton J, Burns JM. The physical performance test predicts aerobic capacity sufficient for independence in early-stage Alzheimer disease. *J Geriatr Phys Ther*. 2012;35:72-8.
- (10) Coelho FG, Stella F, de Andrade LP, Barbieri FA, Santos-Galduroz RF, Gobbi S, et al. Gait and risk of falls associated with frontal cognitive functions at different stages of Alzheimer's disease. *Neuropsychology, development, and cognition Section B, Aging, neuropsychology and cognition*. 2012;19:644-56.
- (11) Muir SW, Speechley M, Wells J, Borrie M, Gopaul K, Montero-Odasso M. Gait assessment in mild cognitive impairment and Alzheimer's disease: the effect of dual-task challenges across the cognitive spectrum. *Gait & posture*. 2012;35:96-100.
- (12) Muir SW, Beauchet O, Montero-Odasso M, Annweiler C, Fantino B, Speechley M. Association of executive function impairment, history of falls and physical performance in older adults: a cross-sectional population-based study in eastern France. *The journal of nutrition, health & aging*. 2013;17:661-5.
- (13) Amboni M, Barone P, Hausdorff JM. Cognitive contributions to gait and falls: evidence and implications. *Movement disorders : official journal of the Movement Disorder Society*. 2013;28:1520-33.
- (14) Forbes D, Forbes SC, Blake CM, Thiessen EJ, Forbes S. Exercise programs for people with dementia. *The Cochrane database of systematic reviews*. 2015;4:CD006489.
- (15) Blankevoort CG, van Heuvelen MJ, Boersma F, Luning H, de Jong J, Scherder EJ. Review of effects of physical activity on strength, balance, mobility and ADL performance in elderly subjects with dementia. *Dement Geriatr Cogn Disord*. 2010;30:392-402.
- (16) Littbrand H, Stenvall M, Rosendahl E. Applicability and effects of physical exercise on physical and cognitive functions and activities of daily living among people with dementia: a systematic review. *American journal of physical medicine & rehabilitation / Association of Academic Physiatrists*. 2011;90:495-518.
- (17) Sobol NA, Hoffmann K, Frederiksen KS, Vogel A, Vestergaard K, Braendgaard H, et al. Effect of aerobic exercise on physical performance in patients with Alzheimer's disease. *Alzheimer's & dementia : the journal of the Alzheimer's Association*. 2016.
- (18) Brett L, Traynor V, Stapley P. Effects of Physical Exercise on Health and Well-Being of Individuals Living With a Dementia in Nursing Homes: A Systematic Review. *J Am Med Dir Assoc*. 2016;17:104-16.
- (19) Laver K, Dyer S, Whitehead C, Clemson L, Crotty M. Interventions to delay functional decline in people with dementia: a systematic review of systematic reviews. *BMJ Open*. 2016;6:e010767.

- 
- (20) Pitkala K, Savikko N, Poysti M, Strandberg T, Laakkonen ML. Efficacy of physical exercise intervention on mobility and physical functioning in older people with dementia: a systematic review. *Exp Gerontol*. 2013;48:85-93.
- (21) Telenius EW, Engedal K, Bergland A. Effect of a high-intensity exercise program on physical function and mental health in nursing home residents with dementia: an assessor blinded randomized controlled trial. *PLoS one*. 2015;10:e0126102.
- (22) Cancela JM, Ayan C, Varela S, Seijo M. Effects of a long-term aerobic exercise intervention on institutionalized patients with dementia. *J Sci Med Sport*. 2015.
- (23) Rao AK, Chou A, Bursley B, Smulofsky J, Jezequel J. Systematic review of the effects of exercise on activities of daily living in people with Alzheimer's disease. *The American journal of occupational therapy : official publication of the American Occupational Therapy Association*. 2014;68:50-6.
- (24) Toots A, Littbrand H, Lindelof N, Wiklund R, Holmberg H, Nordstrom P, et al. Effects of a High-Intensity Functional Exercise Program on Dependence in Activities of Daily Living and Balance in Older Adults with Dementia. *J Am Geriatr Soc*. 2016;64:55-64.
- (25) Burton E, Cavalheri V, Adams R, Browne CO, Boverly-Spencer P, Fenton AM, et al. Effectiveness of exercise programs to reduce falls in older people with dementia living in the community: a systematic review and meta-analysis. *Clinical interventions in aging*. 2015;10:421-34.
- (26) Perrochon A, Tchalla AE, Bonis J, Perucaud F, Mandigout S. Effects of a Multicomponent Exercise Program on Spatio-temporal Gait Parameters, Risk of Falling and Physical Activity in Dementia Patients. *Dementia and geriatric cognitive disorders extra*. 2015;5:350-60.
- (27) Chan WC, Fai Yeung JW, Man Wong CS, Wa Lam LC, Chung KF, Hay Luk JK, et al. Efficacy of physical exercise in preventing falls in older adults with cognitive impairment: a systematic review and meta-analysis. *J Am Med Dir Assoc*. 2015;16:149-54.
- (28) Schwenk M, Zieschang T, Oster P, Hauer K. Dual-task performances can be improved in patients with dementia: a randomized controlled trial. *Neurology*. 2010;74:1961-8.
- (29) Coelho FG, Andrade LP, Pedroso RV, Santos-Galduroz RF, Gobbi S, Costa JL, et al. Multimodal exercise intervention improves frontal cognitive functions and gait in Alzheimer's disease: a controlled trial. *Geriatrics & gerontology international*. 2013;13:198-203.
- (30) Ohman H, Savikko N, Strandberg TE, Pitkala KH. Effect of physical exercise on cognitive performance in older adults with mild cognitive impairment or dementia: a systematic review. *Dement Geriatr Cogn Disord*. 2014;38:347-65.
- (31) Phillips C, Baktir MA, Das D, Lin B, Salehi A. The Link Between Physical Activity and Cognitive Dysfunction in Alzheimer Disease. *Physical therapy*. 2015;95:1046-60.
- (32) Hoffmann K, Sobol NA, Frederiksen KS, Beyer N, Vogel A, Vestergaard K, et al. Moderate-to-High Intensity Physical Exercise in Patients with Alzheimer's Disease: A Randomized Controlled Trial. *J Alzheimers Dis*. 2015.
- (33) Brown BM, Peiffer JJ, Martins RN. Multiple effects of physical activity on molecular and cognitive signs of brain aging: can exercise slow neurodegeneration and delay Alzheimer's disease? *Molecular psychiatry*. 2013;18:864-74.
- (34) Thune-Boyle IC, Iliffe S, Cerga-Pashoja A, Lowery D, Warner J. The effect of exercise on behavioral and psychological symptoms of dementia: towards a research agenda. *Int Psychogeriatr*. 2012;24:1046-57.
- (35) Zhu XC, Yu Y, Wang HF, Jiang T, Cao L, Wang C, et al. Physiotherapy intervention in Alzheimer's disease: systematic review and meta-analysis. *J Alzheimers Dis*. 2015;44:163-74.
- (36) Moyle W, Murfield JE, O'Dwyer S, Van Wyk S. The effect of massage on agitated behaviours in older people with dementia: a literature review. *J Clin Nurs*. 2013;22:601-10.
- (37) Strom BS, Ytrehus S, Grov EK. Sensory stimulation for persons with dementia: a review of the literature. *J Clin Nurs*. 2016;25:1805-34.
- (38) Viggo Hansen N, Jorgensen T, Ortenblad L. Massage and touch for dementia. *The Cochrane database of systematic reviews*. 2006:CD004989.