

# Fysioterapi i behandlingen af type 2-diabetes

## Fuld referenceliste

- (1) Stratton IM, Adler AI, Neil HA, Matthews DR, Manley SE, Cull CA, et al. Association of glycaemia with macrovascular and microvascular complications of type 2 diabetes (UKPDS 35): prospective observational study. *BMJ* 2000 Aug 12;321(7258):405-412.
- (2) Park SW, Goodpaster BH, Strotmeyer ES, Kuller LH, Broudeau R, Kammerer C, et al. Accelerated loss of skeletal muscle strength in older adults with type 2 diabetes: the health, aging, and body composition study. *Diabetes Care* 2007 Jun;30(6):1507-1512.
- (3) Molsted S, Tribler J, Snorgaard O. Musculoskeletal pain in patients with type 2 diabetes. *Diabetes Res Clin Pract* 2012 May;96(2):135-140.
- (4) Umpierre D, Ribeiro PA, Kramer CK, Leitao CB, Zucatti AT, Azevedo MJ, et al. Physical activity advice only or structured exercise training and association with HbA1c levels in type 2 diabetes: a systematic review and meta-analysis. *JAMA* 2011 May 4;305(17):1790-1799.
- (5) Chudyk A, Petrella RJ. Effects of exercise on cardiovascular risk factors in type 2 diabetes: a meta-analysis. *Diabetes Care* 2011 May;34(5):1228-1237.
- (6) Snowling NJ, Hopkins WG. Effects of different modes of exercise training on glucose control and risk factors for complications in type 2 diabetic patients: a meta-analysis. *Diabetes Care* 2006 Nov;29(11):2518-2527.
- (7) Thomas DE, Elliott EJ, Naughton GA. Exercise for type 2 diabetes mellitus. *Cochrane Database Syst Rev* 2006 Jul 19;3:CD002968.
- (8) Bacchi E, Negri C, Zanolin ME, Milanese C, Faccioli N, Trombetta M, et al. Metabolic effects of aerobic training and resistance training in type 2 diabetic subjects: a randomized controlled trial (the RAED2 study). *Diabetes Care* 2012 Apr;35(4):676-682.
- (9) Belli T, Ribeiro LF, Ackermann MA, Baldissera V, Gobatto CA, Galdino da Silva R. Effects of 12-week overground walking training at ventilatory threshold velocity in type 2 diabetic women. *Diabetes Res Clin Pract* 2011 Sep;93(3):337-343.
- (10) Negri C, Bacchi E, Morgante S, Soave D, Marques A, Menghini E, et al. Supervised walking groups to increase physical activity in type 2 diabetic patients. *Diabetes Care* 2010 Nov;33(11):2333-2335.
- (11) Kodama S, Tanaka S, Heianza Y, Fujihara K, Horikawa C, Shimano H, et al. Association Between Physical Activity and Risk of All-Cause Mortality and Cardiovascular Disease in Patients With Diabetes: A meta-analysis. *Diabetes Care* 2013 Feb;36(2):471-479.
- (12) Lebedz-Odrobina D, Kay J. Rheumatic manifestations of diabetes mellitus. *Rheum Dis Clin North Am* 2010 Nov;36(4):681-699.
- (13) Banerjee C, Moon YP, Paik MC, Rundek T, Mora-McLaughlin C, Vieira JR, et al. Duration of diabetes and risk of ischemic stroke: the Northern Manhattan Study. *Stroke* 2012 May;43(5):1212-1217.
- (14) Ehrlich SF, Quesenberry CP, Jr, Van Den Eeden SK, Shan J, Ferrara A. Patients diagnosed with diabetes are at increased risk for asthma, chronic obstructive pulmonary disease, pulmonary fibrosis, and pneumonia but not lung cancer. *Diabetes Care* 2010 Jan;33(1):55-60.
- (15) Karstoft K, Wallis GA, Pedersen BK, Solomon TP. The effects of interval- vs. continuous exercise on excess post-exercise oxygen consumption and substrate oxidation rates in subjects with type 2 diabetes. *Metabolism* 2016 Sep;65(9):1316-25.
- (16) Coyle D, Coyle K, Kenny GP, Boule NG, Wells GA, Fortier M, et al. Cost-effectiveness of exercise programs in type 2 diabetes. *Int J Technol Assess Health Care* 2012 Jul;28(3):228-234.
- (17) Roessler K. Fysisk aktivitet som behandling. Odense: Syddansk Universitetsforlag; 2007.
- (18) Sundhedsstyrelsen. Resultatopsamling af motion på recept i Danmark. Version: 2,0 ed. København: Sundhedsstyrelsen; 2010.
- (19) Deakin T, McShane CE, Cade JE, Williams RD. Group based training for self-management strategies in people with type 2 diabetes mellitus. *Cochrane Database Syst Rev* 2005 Apr 18;(2):CD003417.
- (20) Molsted S, Tribler J, Poulsen PB, Snorgaard O. The effects and costs of a group-based education programme for self-management of patients with Type 2 diabetes. A community-based study. *Health Educ Res* 2012 Oct;27(5):804-813.
- (21) Diabetes - Statens Serum Institut Available at: <http://www.ssi.dk/Sundhedsdataogit/Dataformidling/Sundhedsdata/Diabetes.aspx>. Accessed 2/20/2013, 2013.