


Fysioterapi til behandling af mennesker med skizofreni

Fuld referenceliste

- (1) L Nyboe BH, ed. *Lærebog i psykiatri for ergoterapeuter og fysioterapeuter*. 1.th ed. København: Munksgaards Forlag; 2009.
- (2) Nordentoft M, Ahlstand L, Christensen A, et al. *Referenceprogram for skizofreni*. København: Sundhedsstyrelsen; 2004.
- (3) Jørgensen P, ed. *Psykiske sygdomme og kroppen*. Kap. 4 1.udgave ed. København: Munksgaard Danmark; 2011.
- (4) Stanghellini G, Ballerini M, Blasi S, et al. *The bodily self: A qualitative study of abnormal bodily phenomena in persons with schizophrenia*. *Compr Psychiatry*. 2014;55(7):1703-1711. doi: 10.1016/j.comppsy.2014.06.013.
- (5) Stanghellini G. *Embodiment and schizophrenia*. *World Psychiatry*. 2009;8(1):56-59.
- (6) Roxendal G. *Body awareness therapy and the body awareness scale, treatment and evaluation in psychiatric physiotherapy*. Kaallerud: Kompendietrykkeriet; 1985.
- (7) Hedlund L, Gyllensten AL. *The experiences of basic body awareness therapy in patients with schizophrenia*. *J Bodyw Mov Ther*. 2010;14(3):245-254. doi: 10.1016/j.jbmt.2009.03.002.
- (8) Vancampfort D, Probst M, Scheewe T, et al. *Relationships between physical fitness, physical activity, smoking and metabolic and mental health parameters in people with schizophrenia*. *Psychiatry Res*. 2013;207(1-2):25-32. doi: dx.doi.org/10.1016/j.psychres.2012.09.026.
- (9) Nyboe L, Vestergaard CH, Moeller MK, Lund H, Videbech P. *Metabolic syndrome and aerobic fitness in patients with first-episode schizophrenia, including a 1-year follow-up*. *Schizophr Res*. 2015;168(1-2):381-387. doi: 10.1016/j.schres.2015.07.053 [doi].
- (10) Sundhedsstyrelsen. *Fysisk aktivitet - håndbog om forebyggelse og behandling*. 2011.
- (11) Soundy A, Stubbs B, Probst M, Hemmings L, Vancampfort D. *Barriers to and facilitators of physical activity among persons with schizophrenia: A survey of physical therapists*. *PS*. 2014;65(5):693-696. dx.doi.org/10.1176/appi.ps.201300276. doi: 10.1176/appi.ps.201300276.
- (12) Vancampfort D, De Hert M, Stubbs B, et al. *Negative symptoms are associated with lower autonomous motivation towards physical activity in people with schizophrenia*. *Compr Psychiatry*. 2015;56:128-132.
- (13) Vancampfort D, Knapen J, Probst M, Scheewe T, Remans S, De Hert M. *A systematic review of correlates of physical activity in patients with schizophrenia*. *Acta Psychiatr Scand*. 2012;125(5):352-362.
- (14) Bassilios B, Judd F, Pattison P. *Why don't people diagnosed with schizophrenia spectrum disorders (SSDs) get enough exercise?* *Australas Psychiatry*. 2013. doi: 10.1177/1039856213510575.
- (15) Vancampfort D, Wampers M, Mitchell AJ, et al. *A meta-analysis of cardio-metabolic abnormalities in drug naive, first-episode and multi-episode patients with schizophrenia versus general population controls*. *World Psychiatry*. 2013;12(3):240-250. doi: 10.1002/wps.20069
- (16) Mitchell AJ, Vancampfort D, De Hert M, Yu W, De Hert M. *Is the prevalence of metabolic syndrome and metabolic abnormalities increased in early schizophrenia? A comparative meta-analysis of first episode, untreated and treated patients*. *Schizophr Bull*. 2013;39(2):295-305. doi: 10.1093/schbul/sbs082
- (17) De Hert M, Dekker JM, Wood D, Kahl KG, Holt RI, Moller HJ. *Cardiovascular disease and diabetes in people with severe mental illness position statement from the european psychiatric association (EPA), supported by the european association for the study of diabetes (EASD) and the european society of cardiology (ESC)*. *Eur Psychiatry*. 2009;24(6):412-424.
- (18) De Hert M, Vancampfort D, Correll CU, et al. *Guidelines for screening and monitoring of cardiometabolic risk in schizophrenia: Systematic evaluation*. *Br J Psychiatry*. 2011;199(2):99-105.
- (19) Huang Y, Liu X. *Leisure-time physical activity and the risk of metabolic syndrome: Meta-analysis*. *Eur J Med Res*. 2014;19:22-783X-19-22. doi: 10.1186/2047-783X-19-22.
- (20) Vancampfort D, De Hert M, Skjerven LH, et al. *International organization of physical therapy in mental health consensus on physical activity within multidisciplinary rehabilitation programmes for minimising cardio-metabolic risk in patients with schizophrenia*. *Disabil Rehabil*. 2012;34(1):1-12. doi: 10.3109/09638288.2011.587090.
- (21) Laursen TM, Nordentoft M, Mortensen PB. *Excess early mortality in schizophrenia*. *Annu Rev Clin Psychol*. 2014;10:425-448. doi: 10.1146/annurev-clinpsy-032813-153657.
- (22) Moltke A. *Kroppens ambivalens*. Copenhagen: Frydenlund; 2010.



(23) Sundhedsstyrelsen. Sygdomsbyrden i danmark- sygdomme. 2015.

(24) De Hert M, Correll CU, Bobes J, et al. Physical illness in patients with severe mental disorders. I. prevalence, impact of medications and disparities in health care. *World Psychiatry*. 2011;10(1):52-77.

(25) Vancampfort D, Stubbs B, Ward PB, Teasdale S, Rosenbaum S. Why moving more should be promoted for severe mental illness. *Lancet Psychiatry*. 2015;2(4):295-0366(15)00099-1. Epub 2015 Mar 31. doi: 10.1016/S2215-0366(15)00099-1.