

Stillesiddende adfærd – også en helbredsrisiko hos personer med funktionsbegrænsning?

Af Mette Aadahl og Peter Gjerndrup Aagaard
Fysioterapeuten nr. 4 2013.

Litteratur

- (1) Overgaard K, Grøntved A, Nielsen K, Dahl-Petersen IK, Aadahl M. Stillesiddende adfærd - en helbredsrisiko? København: Vidensråd for Forebyggelse; 2012: 1-112.
- (2) Biddle S, Cavill N, Ekelund U, Gorely T, Griffiths M, Jago R, et al. Sedentary Behaviour and Obesity: Review of the Current Scientific Evidence. London: Department of Health, UK; 2010: 1-126.
- (3) Schofield G, Quigley R, Brown R. Does sedentary behaviour contribute to chronic disease or chronic disease risk in adults? New Zealand: Committee of Agencies for Nutrition Action; 2009: 1-128.
- (4) Kiens B et al. Fysisk inaktivitet - konsekvenser og sammenhænge. Søborg: Motions- og Ernæringsrådet; 2007: 1-148.
- (5) Sedentary Behaviour Research Network. Letter to the editor: standardized use of the terms "sedentary" and "sedentary behaviours". *Appl Physiol Nutr Metab* 2012;37(3):540-2.
- (6) Pate RR, O'Neill JR, Lobelo F. The evolving definition of "sedentary". *Exerc Sport Sci Rev* 2008;36(4):173-8.
- (7) Fysisk aktivitet - håndbog om forebyggelse og behandling. København: Sundhedsstyrelsen, 2011.
- (8) Matthiesen J et al. Fysisk aktivitet i den voksne danske befolkning 2003-2006. Søborg: DTU Fødevareinstituttet; 2009.
- (9) Statistisk Årbog 2010. København: Danmarks Statistik, 2010.
- (10) Hammer-Helmich L, Buhelt LP, Andreasen AH, Robinson KM, Hilding-Nørkjær H, Glümer C. Region Hovedstaden. Sundhedsprofil for regioner og kommuner 2010. Glostrup: Forskningscenter for Forebyggelse og Sundhed, 2010.
- (11) Ortiz B. Sundhedsprofil Region Sjælland 2010. Sorø: Region Sjælland, Kvalitet og Udvikling 2010.
- (12) Christensen AI, Ekholm O, Davidsen M, et al. Sundhed & Sygelighed i Danmark 2010 - og udviklingen siden 1987. København: Statens Institut for Folkesundhed, Syddansk Universitet, 2012.
- (13) Grontved A, Hu FB. Television viewing and risk of type 2 diabetes, cardiovascular disease, and all-cause mortality: a meta-analysis. *JAMA* 2011;305(23):2448-55.
- (14) Thorp AA, Owen N, Neuhaus M et al. Sedentary behaviors and subsequent health outcomes in adults a systematic review of longitudinal studies, 1996-2011. *Am J Prev Med* 2011;41(2):207-15.
- (15) Tremblay MS, Colley RC, Saunders TJ et al. Physiological and health implications of a sedentary lifestyle. *Appl Physiol Nutr Metab* 2010;35(6):725-40.
- (16) Dunstan DW, Barr EL, Healy GN et al. Television viewing time and mortality: the Australian Diabetes, Obesity and Lifestyle Study (AusDiab). *Circulation* 2010;121(3):384-91.

- (17) Ford ES, Schulze MB, Kroger J et al. Television watching and incident diabetes: Findings from the European Prospective Investigation into Cancer and Nutrition-Potsdam Study. *J Diabetes* 2010;2(1):23-7.
- (18) Katzmarzyk PT, Church TS, Craig CL et al. Sitting time and mortality from all causes, cardiovascular disease, and cancer. *Med Sci Sports Exerc* 2009;41(5):998-1005.
- (19) Stamatakis E, Hamer M, Dunstan DW. Screen-based entertainment time, all-cause mortality, and cardiovascular events: population-based study with ongoing mortality and hospital events follow-up. *J Am Coll Cardiol* 2011;57(3):292-9.
- (20) Wijndaele K, Brage S, Besson H et al. Television viewing and incident cardiovascular disease: prospective associations and mediation analysis in the EPIC Norfolk Study. *PLoS One* 2011;6(5):e20058.
- (21) van Uffelen JG, Wong J, Chau JY et al. Occupational sitting and health risks: a systematic review. *Am J Prev Med* 2010;39(4):379-88.
- (22) Lynch BM. Sedentary behavior and cancer: a systematic review of the literature and proposed biological mechanisms. *Cancer Epidemiol Biomarkers Prev* 2010;19(11):2691-709.
- (23) Healy GN, Dunstan DW, Salmon J et al. Breaks in sedentary time: beneficial associations with metabolic risk. *Diabetes Care* 2008;31(4):661-6.
- (24) World Health Organization/ World Bank. World Report on Disability. World Health Organization: 2011.
- (25) Rand D, Eng JJ, Tang PF et al. How active are people with stroke?: use of accelerometers to assess physical activity. *Stroke* 2009;40(1):163-8.
- (26) Cavanaugh JT, Coleman KL, Gaines JM et al. Using step activity monitoring to characterize ambulatory activity in community-dwelling older adults. *J Am Geriatr Soc* 2007;55(1):120-4.
- (27) Chastin SF, Baker K, Jones D et al. The pattern of habitual sedentary behavior is different in advanced Parkinson's disease. *Mov Disord* 2010;25(13):2114-20.
- (28) Healy GN, Wijndaele K, Dunstan DW et al. Objectively measured sedentary time, physical activity, and metabolic risk: the Australian Diabetes, Obesity and Lifestyle Study (AusDiab). *Diabetes Care* 2008;31(2):369-71.
- (29) Matthews CE, Chen KY, Freedson PS et al. Amount of time spent in sedentary behaviors in the United States, 2003-2004. *Am J Epidemiol* 2008;167(7):875-81.
- (30) Hagstromer M, Oja P, Sjostrom M. Physical activity and inactivity in an adult population assessed by accelerometry. *Med Sci Sports Exerc* 2007;39(9):1502-8.
- (31) Manns PJ, Dunstan DW, Owen N et al. Addressing the nonexercise part of the activity continuum: a more realistic and achievable approach to activity programming for adults with mobility disability? *Phys Ther* 2012;92(4):614-25.