

Referenceliste til artiklen

Tung styrketræning til patienter med brystkræft

AF FYSIOTERAPEUT, PH.D. KIRA BRISSON BLOOMQUIST
FYSIOTERAPEUTEN NR. 6 2018

1. DiSipio T, Rye S, Newman B, Hayes S. **Incidence of unilateral arm lymphoedema after breast cancer: a systematic review and meta-analysis.** *Lancet Oncol.* 2013;14(6):500-15.
2. Fu MR, Ridner SH, Hu SH, Stewart BR, Cormier JN, Armer JM. **Psychosocial impact of lymphedema: a systematic review of literature from 2004 to 2011.** *Psychooncology.* 2013;22(7):1466-84.
3. Boyages J, Kalfa S, Xu Y, Koelmeyer L, Mackie H, Viveros H, et al. **Worse and worse off: the impact of lymphedema on work and career after breast cancer.** *Springerplus.* 2016;5:657.
4. Paramanandam VS, Roberts D. **Weight training is not harmful for women with breast cancer-related lymphoedema: a systematic review.** *J Physiother.* 2014;60(3):136-43.
5. Cheema BS, Kilbreath SL, Fahey PP, Delaney GP, Atlantis E. **Safety and efficacy of progressive resistance training in breast cancer: a systematic review and meta-analysis.** *Breast Cancer Res Treat.* 2014;148(2):249-68.
6. Schmitz KH. **Balancing lymphedema risk: exercise versus deconditioning for breast cancer survivors.** *Exerc Sport Sci Rev.* 2010;38:17-24.
7. Nelson NL. **Breast Cancer-Related Lymphedema and Resistance Exercise: A Systematic Review.** *J Strength Cond Res.* 2016;30(9):2656-65.
8. Csapo R, Alegre LM. **Effects of resistance training with moderate vs heavy loads on muscle mass and strength in the elderly: A meta-analysis.** *Scand J Med Sci Sports.* 2016;26(9):995-1006.
9. Donnelly JE, Blair SN, Jakicic JM, Manore MM, Rankin JW, Smith BK. **American College of Sports Medicine Position Stand. Appropriate physical activity intervention strategies for weight loss and prevention of weight regain for adults.** *Med Sci Sports Exerc.* 2009;41(2):459-71.
10. Kohrt WM, Bloomfield SA, Little KD, Nelson ME, Yingling VR. **American College of Sports Medicine Position Stand: physical activity and bone health.** *Med Sci Sports Exerc.* 2004;36(11):1985-96.
11. Bloomquist K, Karlsmark T, Christensen KB, Adamsen L. **Heavy resistance training and lymphedema: prevalence of breast cancer-related lymphedema in participants of an exercise intervention utilizing heavy load resistance training.** *Acta Oncol.* 2014;53(2):216-25.
12. Adamsen L, Quist M, Andersen C, Moller T, Herrstedt J, Kronborg D, et al. **Effect of a multimodal high intensity exercise intervention in cancer patients undergoing chemotherapy: randomised controlled trial.** *BMJ.* 2009;339:b3410.

13. Bloomquist K, Hayes S, Adamsen L, Moller T, Christensen KB, Ejlertsen B, et al. **A randomized cross-over trial to detect differences in arm volume after low- and heavy-load resistance exercise among patients receiving adjuvant chemotherapy for breast cancer at risk for arm lymphedema: study protocol.** BMC Cancer. 2016;16:517.
14. Bloomquist K, Oturai P, Steele ML, Adamsen L, Moller T, Christensen KB, et al. **Heavy-Load Lifting: Acute Response in Breast Cancer Survivors at Risk for Lymphedema.** Med Sci Sports Exerc. 2018;50(2):187-95.
15. Moller T, Lillelund C, Andersen C, Ejlertsen B, Norgaard L, Christensen KB, et al. **At cancer diagnosis: a 'window of opportunity' for behavioural change towards physical activity. A randomised feasibility study in patients with colon and breast cancer.** BMJ Open. 2013;3(11):e003556.